Author Index

Aboul-Enein, H.Y.
—— and Islam, M. R.
Direct separation and optimization of timolol enantiomers on a cellulose tris-3,5-dimethylphenylcarbamate high-performance liquid chromatographic chiral stationary phase 109
Antonozzi, I., see Moretti, F. 131
Apps, P. J.
High-precision sampling of sub-nanogram, low-parts-per-billion solutes from liquids using the dynamic solvent effect 271
Asakawa, N., see Miwa, T. 89
Belliardo, F., see Lucarelli, C. 167
Betto, P., see Lucarelli, C. 167
Betts, T. J.
Chromatographic evaluation of boldine and associated alkaloids in Boldo 373
Bianco, O., see Petrarulo, M. 223
Bilic, N.
—— and Sieber, R.
Determination of flavins in dairy products by high-performance liquid chromatography using sorboflavin as internal standard 359
Birarelli, M., see Moretti, F. 131
Bonora, A., see Menziani, E. 396
Breyer, E. D., see Strasters, J. K. 17
Briggs, L. M., see Treado, P. J. 341
Brinkman, U. A. Th., see Hogendoorn, E. A. 243
——, see Kwakman, P. J.M. 155
Burgers, I., see Hordijk, C. A. 317
Cappenberg, T. E., see Hordijk, C. A. 317
Carducci, C., see Moretti, F. 131
Carner, R. C., see Carvajal, M. 379
Carvajal, M.
——, Mulholland, F. and Carner R. C.
Comparison of the EASI-EXTRACT immunoaffinity concentration procedure with the AOAC CB method for the extraction and quantitation of aflatoxin B_1 in raw ground unskinned peanuts 379
Corradini, C., see Lucarelli, C. 167
Covey, T. R., see Huang, E. 257
Cox, G. B., see Snyder, L. R. 404
Csató, E.
——, Fülöp, N. and Szabó, G.
Preparation and comparison of a pentafluorophenyl stationary phase for reversed-phase liquid chromatography 79
De Jong, A. P. J. M., see Hogendoorn, E. A. 243
De Jong, G. J., see Kwakman, P. J. M. 155
Demchak, R. J.
—— and MacConnell, J. G.
Normal-phase chromatography and post-column colorimetric detection of abamectin-8,9-oxide 353
Dufresne, M., see Jiménez, J. 333
Dziwiński, E., see Szymanowski, J. 325
El Fallah, M. Z.
——, Golshan-Shirazi, S. and Guiochon, G.
Theoretical study of the effect of a difference in column saturation capacities for the two components of a binary mixture on their elution band profiles and separation in non-linear chromatography I
Etcheverry, M. F., see Pierron, C. 367
Fallah, M. Z. el, see El Fallah, M. Z. I
Fidge, N., see Tetaz, T. 147
Fourmy, D., see Jiménez, J. 333
Fu, R., see Zhu, P. 390
Fukuda, T.
——, Kohara, N., Onogi, Y. and Inagaki, H.
Separation of simple ions by gel chromatography. I. Simple model of separation for single-salt systems 59
Fülöp, N., see Csató, E. 79
Galushko, S.V.
——, Shishkina, I. P., Soloshonok, V. A. and Kukhar, V. P.
Ligand-exchange chromatography of α-trifluoromethyl-α-amino acids on chiral sorbents 115
Gehas, J.
—— and Wetlaufer, D. B.
Isocratic hydrophobic interaction chromatography of dansyl amino acids. Correlation of hydrophobicity and retention parameters 123
Giambenedetti, M., see Lucarelli, C. 167
Golshan-Shirazi, S.
—— and Guiochon, G.
Influence of pressure on the maximum production rate in preparative liquid chromatography 402
——, see El Fallah, M. Z. I
Gordon, S. M.
Identification of exposure markers in smokers' breath 291
Greef, J. van der, see Kokkonen, P. 35
——, see Stegehuis, D. S. 137
Grego, B., see Tetaz, T. 147
Grönberg, L., see Gyllenhaal, O. 303
Guiochon, G., see El Fallah, M. Z. I
——, see Golshan-Shirazi, S. 402

Gyllenhaal, O.
——, Grönberg, L. and Vessman, J.
Determination of hydrazine in hydralazine by capillary gas chromatography with nitrogen-selective detection after benzaldehyde derivatization 303
Hatano, H., see Naikwadi, K. P. 281
Henion, J., see Huang, E. 257
Hogendoorn, E. A.
——, De Jong, A. P. J. M., Van Zoonen, P. and Brinkman, U. A. Th.
Reversed-phase liquid chromatographic column switching for the trace-level determination of polar compounds. Application to chloroallyl alcohol in ground water 243
Hordijk, C. A.
——, Burgers, I., Phylipsen, G. J. M. and Cappenberg, T. E.
Trace determination of lower volatile fatty acids in sediments by gas chromatography with chemically bonded FFAP columns 317
Huang, E.
——, Henion, J. and Covey, T. R.
Packed-column supercritical fluid chromatography–mass spectrometry and supercritical fluid chromatography–tandem mass spectrometry with ionization at atmospheric pressure 257
Inagaki, H., see Fukuda, T. 59
Islam, M. R., see Aboul-Enein, H. Y. 109
Jarrett, H. W., see Lee, H. G. 69
Jiménez, J.
——, Dufresne, M., Poirot, S., Vaysse, N. and Fourmy, D.
Electric properties of photoaffinity-labelled pancreatic A-subtype cholecystokinin 333
Jong, A. P. J. M. de, see Hogendoorn, E. A. 243
Jong, G. J. de, see Kwakman, P. J. M. 155
Kanhere, S. R.
—— and Scott, P. M.
Heptafluorobutyrylation of trichothecenes using a solid-phase catalyst 384
Karasek, F. W., see Naikwadi, K. P. 281
Kecorius, E., see Tetaz, T. 147
Kemmelmeier, C., see Paterson, R. R. M. 195
Khaledi, M. G., see Strasters, J. K. 17
Koelewijn, H., see Kwakman, P. J. M. 155
Kohara, N., see Fukuda, T. 59
Kokkonen, P.
——, Schröder, E., Niessen, W. M. A., Tjaden, U. R. and Van der Greef J.
Important parameters in liquid chromatography–continuous flow fast atom bombardment mass spectrometry 35
Kool, I., see Kwakman, P. J. M. 155
Kukhar, V. P., see Galushko, S. V. 115
Kuroda, H., see Miwa, T. 89
Kusz, P., see Szymanowski, J. 325
Kwakman, P. J. M.
——, Koelewijn, H., Kool, I., Brinkman, U. A. Th. and De Jong, G. J.
Naphthalene- and anthracene-2,3-dialdehyde as precolumn labelling reagents for primary amines using reversed- and normal-phase liquid chromatography with peroxyoxalate chemiluminescence detection 155
Ledouble, G., see Pierron, C. 367
Lee, H. G.
—— and Jarrett, H. W.
Glucose-silica, an improved medium for high-pressure gel filtration chromatography 69
Linari, F., see Petrarulo, M. 223
Liu, H., see Zhu, P. 390
Lodi, G., see Menziani, E. 396
Lucarelli, C.
——, Betto, P., Ricciarello, G., Giambenedetti, M., Corradini, C., Stocchi, F. and Belliardo, F.
Simultaneous measurement of L-DOPA, its metabolites and carbidopa in plasma of Parkinsonian patients by improved sample pretreatment and high-performance liquid chromatographic determination 167
MacConnell, J. G., see Demchak, R. J. 353
Marangella, M., see Petrarulo, M. 223
Mentasti, E., see Petrarulo, M. 223
Menziani, E.
——, Tosi, B., Bonora, A., Reschiglian, P. and Lodi, G.
Automated multiple development high-performance thin-layer chromatographic analysis of natural phenolic compounds 396
Miller, L.
—— and Weyker, C.
Analytical and preparative resolution of enantiomers of prostaglandin precursors and prostaglandins by liquid chromatography on derivatized cellulose chiral stationary phases 97
Minocha, R., see Minocha, S. C. 177
Minocha, S. C.
——, Minocha, R. and Robie, C.A.
High-performance liquid chromatographic method for the determination of dansyl-polyamines 177
Miwa, T.
——, Kuroda, H., Sakashita, S., Asakawa, N. and Miyake, Y.
Characteristics of ovomucoid-conjugated columns in the direct liquid chromatographic resolution of racemic compounds 89
Miyake, Y., see Miwa, T. 89

Moretti, F.
——, Birarelli, M., Carducci, C., Pontecorvi, A. and Antonozzi, I.
Simultaneous high-performance liquid chromatographic determination of amino acids in a dried blood spot as a neonatal screening test 131
Morris, M. D., see Treado, P. J. 341
Mulholland, F., see Carvajal, M. 379
Naikwadi, K. P.
——, Karasek, F. W. and Hatano, H.
Analyses of polychlorinated dibenzo-*p*-dioxins and dibenzofurans and precursors in fly ash samples collected at different point in post-combustion zone of Japanese machida incinerator 281
Niessen, W. M. A., see Kokkonen, P. 35
Onogi, Y., see Fukuda, T. 59
Panas, J. M., see Pierron, C. 367
Parkin, J. E.
High-performance liquid chromatographic investigation of the interaction of phenylmercuric nitrate and sodium metabisulphite in eye drop formulations 233
Paterson, R. R. M.
——, and Kemmelmeier, C.
Neutral, alkaline and difference ultraviolet spectra of secondary metabolites from *Penicillium* and other fungi, and comparisons to published maxima from gradient high-performance liquid chromatography with diode-array detection 195
Pellegrino, S., see Petrarulo, M. 223
Petrarulo, M.
——, Bianco, O., Marangella, M., Pellegrino, S., Linari, F and Mentasti, E.
Ion chromatographic determination of plasma oxalate in healthy subjects, in patients with chronic renal failure and in cases of hyperoxaluric syndromes 223
Phylipsen, G. J. M., see Hordijk, C. A. 317
Pierron, C.
——, Panas, J. M. Etcheverry, M. F. and Ledouble, G.
Determination of 10-camphorsulphonates in pharmaceutical formulations by high-performance liquid chromatography 367
Poirot, S., see Jiménez, J. 333
Pontecorvi, A., see Moretti, F. 131
Reschiglian, P., see Menziani, E. 396
Ricciarello, G., see Lucarelli, C. 167
Robie, C. A., see Minocha, S. C. 177
Rodgers, A. H., see Strasters, J. K. 17
Sakashita, S., see Miwa, T. 89
Schröder, E., see Kokkonen, P. 35
Scott, P. M., see Kanhere, S. R. 384
Shi, L., see Zhu, P. 390
Shiskina, I. P., see Galushko, S. V. 155
Sieber, R., see Bilic, N. 359
Skarping, G., see Tiljander, A. 185
Snyder, L. R.
—— and Cox, G. B.
Influence of pressure on the maximum production rate in preparative liquid chromatography. Reply to the letter of S. Golshan-Shirazi and G. Guiochon 404
Soloshonok, V. A., see Galushko, S. V. 115
Stegehuis, D. S.
——, Tjaden, U. R. and Van der Greef, J.
Bioanalysis of the peptide des-enkephalin-γ-endorphin. On-line sample pretreatment using membrane dialysis and solid-phase isolation 137
Stocchi, F., see Lucarelli, C. 167
Strasters, J. K.
——, Breyer, E. D., Rodgers, A. H. and Khaledi, M. G.
Simultaneous optimization of variables influencing selectivity and elution strength in micellar liquid chromatography. Effect of organic modifier and micelle concentration 17
Szabó, G., see Csató, E. 79
Szymanowski, J.
——, Kusz, P. and Dziwiński, E.
Degradation and analysis of commercial polyoxyethylene glycol mono(4-alkylphenyl) ethers 325
Tetaz, T.
——, Kecorius, E., Grego, B. and Fidge, N.
Separation of human apolipoproteins A-IV, A-I and E by reversed-phase high-performance liquid chromatography on a TSK Phenyl-5 PW column 147
Tiljander, A.
—— and Skarping, G.
Determination of 4,4′-methylenedianiline in hydrolysed human urine using liquid chromatography with UV detection and peak identification by absorbance ratio 185
Tjaden, U. R., see Kokkonen, P. 35
——, see Stegehuis, D. S. 137
Tosi, B., see Menziani, E. 396
Treado, P. J.
——, Briggs, L. M. and Morris, M. D.
Hadamard transform photothermal deflection densitometry of electrophoretically blotted proteins 341
Van der Greef, J., see Kokkonen, P. 35
——, see Stegehuis, D. S. 137
Van Zoonen, P., see Hogendoorn, E. A. 243
Vaysse, N., see Jiménez, J. 333
Vessman, J., see Gyllenhaal, O. 303

Vilenchik, L. Z.
Theoretical aspects of the chromatographic analysis of heterogeneous polymers 49
Wetlaufer, D. B., see Gehas, J. 123
Weyker, C., see Miller, L. 97
Ye, M., see Zhu, P. 390
Zhu, P.
——, Ye, M., Shi, L., Liu, H. and Fu, R.
Poly(styrene–dimethylsiloxane) block copolymer as a stationary phase for capillary gas chromatography 390
Zoonen, P. van, see Hogendoorn, E. A. 243

Erratum

J. Chromatogr., 504 (1990) 391–401

Page 391, Summary, third sentence should read

"The individual truxillines were characterized via high-performance liquid chromatography–diode array detection, capillary gas chromatography–electron capture detection, capillary gas chromatography–electron ionization mass spectrometry and capillary supercritical fluid chromatography–flame ionization detection."

Page 395, 4th text line, "tritium" should read "3-H".